



APÉNDICE 4.2: MORTALIDAD NATURAL

Tasas de mortalidad natural adoptadas por el SCRS para las principales especies

Spp.	Stock	M (por año) ¹ por edad											Ref.
		0	1	2	3	4	5	6	7	8	9	10	
ALB	North, South		0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3			Anon. (1990)
BET	Atl.	0.8	0.8	0.4	0.4	0.4	0.4	0.4	0.4				Anon. (1984)
BFT	East		0.49	0.24	0.24	0.24	0.24	0.20	0.175	0.15	0.125	0.10	Anon. (1997);
BFT	West		0.14	0.14	0.14	0.14	0.14	0.14	0.14	0.14	0.14	0.14	?
SKJ	Atl.	0.8	0.8	0.8	0.8								Anon. (1999)
SWO	North		0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	Anon. (1987)
SWO	Med		0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	Anon. (1996)
YFT	Atl.	0.8	0.8	0.6	0.6	0.6	0.6						Anon. (1984)

¹ Last age with an M value is usually a "plus group"

REFERENCIAS

- ANON. 1984. Meeting of the Working Group on Juvenile Tropical Tunas. Collect. Vol. Sci. Pap. ICCAT, 21 (1): 1-289.
- ANON. 1987. Report of the Swordfish Assessment Workshop. Collect. Vol. Sci. Pap. ICCAT, 26 (2): 339-395.
- ANON. 1990. Report of the 1989 ICCAT Albacore Workshop. Collect. Vol. Sci. Pap. ICCAT, 31: 73-210.
- ANON. 1996. Report of the First meeting of the Ad Hoc GFCM-ICCAT Working group on Stocks of Large Pelagic Fishes in the Mediterranean. Collect. Vol. Sci. Pap. ICCAT, 45(1): 1-60.
- ANON. 1997. Report of the 1996 ICCAT SCRS Bluefin Tuna Stock Assessment Session. Collect. Vol. Sci. Pap. ICCAT, 46 (1): 1-186.
- ANON. 1999. 1998 Detailed Report on Skipjack Tuna. Collect. Vol. Sci. Pap. ICCAT, 49 (3): 123-158.